

# **UNIVERSITY OF CALIFORNIA**

## **Northern Regional Library Facility**

### **ELECTRONIC DELIVERY COVER SHEET**

#### **WARNING CONCERNING COPYRIGHT RESTRICTIONS**

**The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted materials.**

**Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.**

**This institution reserves the right to refuse to accept a copying order if, in its judgment, fulfillment of the order would involve violation of copyright law.**

PROCEEDINGS  
OF  
THE GEOLOGICAL SOCIETY  
OF LONDON.

NOVEMBER 1838 to JUNE 1842.

---

VOL. III.

---



LONDON:

PRINTED BY R. AND J. E. TAYLOR, RED LION COURT, FLEET STREET;

AND SOLD AT THE APARTMENTS OF THE SOCIETY,  
SOMERSET HOUSE.

1842.

proposes to designate it by the name of *C. decussata*. The collection of the late Mr. Crow, now in the Canterbury Museum, contains three specimens of this fossil, which, with two others presented by Mr. E. Crow to the Geological Society, are said by the author on the authority of that gentleman, to have been found in digging a ha-ha at Nash Court, about two miles from Faversham. Mr. Trimmer likewise mentions as independent evidence of these fossils having been found at Nash Court, that it is stated on both editions of Mr. Greenough's Map, that siliceous fossils occur there.

A careful examination of the specimens presented to the Geological Society by Mr. E. Crow, has verified the correctness of Mr. Parkinson's opinion, that the Faversham shell is specifically distinct from the *Cucullææ* of the greensands.

The strata of Nash Court, Mr. Trimmer says, undoubtedly belong to the lowest sands of the London clay, and to that portion which is very near the junction with the chalk.

In the village of Boughton, not far from Nash Court, Mr. Trimmer has examined two sections situated to the east and west of the 50th mile-stone, and nearly on a level with the ha-ha, in which the *Cucullæa decussata* was found. The strata consisted of white and ferruginous sand with layers of ferruginous clay, in some parts considerably indurated. He did not observe any organic remains, but shells are reported to have been found in the eastern section. At a greater elevation on the side of Boughton Hill, and a little below the junction of the sands with the brown clay, which forms the summit of the hill, are courses of geodes of a ferruginous sandstone very like some of the Wealden sandstone, and lined with mammillary siliceous deposits and quartz crystals. Casts of plastic clay shells are occasionally found in the sandstone, but more abundantly in the alternate layers of indurated ferruginous clay. From a bed four feet thick of this sandstone worked in a quarry in the wood on the side of Boughton Hill, Mr. Trimmer obtained casts of *Calyptrea trochiformis*, *Rostellaria Sowerbii* (*Strombus pes Pelicani* of Mr. Parkinson), *Potamides intermedium*, and a Venus which has been considered a variety of *V. ovalis*, but is clearly distinct. Similar remains are stated by the author to occur in the upper part of the cliffs at Reculver, either in loose masses, or sand slightly indurated. In conclusion, Mr. Trimmer acknowledges the assistance which he received in preparing the communication.

4. "A description of a portion of the skeleton of the Cetiosaurus, a gigantic extinct Saurian Reptile occurring in the Oolitic formations of different portions of England," by Professor Owen, F.R.S., F.G.S.

The remains described in this memoir consist of vertebrae and bones of the extremities obtained by Mr. Kingdon from the oolite quarries of Chipping Norton, in Oxfordshire; of vertebrae and other bones from the oolite of Blisworth, near Northampton, transmitted to the author by Miss Baker; and of other remains from the oolite of Staple Hill, Wotton, three miles north-west of Woodstock; from the oolite near Buckingham; the Portland stone at Garsington and

Thame, in the collection of Dr. Buckland : Mr. Owen has likewise examined a vertebra and some bones of the extremities of the same saurian from the Yorkshire oolite, and preserved in the Scarborough Museum.

*Caudal Vertebra.*—A caudal vertebra from near Buckingham, which presented the anchylosed neural arch entire, but with the transverse, oblique and spinous processes broken off, equalled in dimensions a middle caudal vertebra of a full-sized whale, the antero-posterior diameter being five inches, the transverse eight inches six lines, and the vertical seven inches. The sides and under part of the centrum are described as very concave ; and the shape of the articular extremities as nearly circular, with a greater concavity in the anterior one than in the posterior. The posterior hæmapophysial articular surfaces slope downwards and forwards in the form of semicircular facets for nearly two inches upon the under surfaces. The neurapophyses commence close to the anterior surface of the centrum, their antero-posterior extent being three and a half inches, and they meet at a rather acute angle above the spinal canal. The vertical diameter of the spinal canal was one inch nine lines, the transverse two inches, and the breadth of the base of the neural arch, from the outside of the neurapophyses, five inches three lines. The transverse process is developed from the centrum just below the neurapophysial suture. In all the caudal vertebrae of the *Cetiosaurus* the posterior half of the centrum is left uncovered by the neural arch.

The substance of another fractured vertebra, showing the upper third of the centrum, presented an uniform coarse spongy texture ; whilst in a third specimen, which exhibited also a similar texture, the layers affected a direction parallel with the articular extremities for about half an inch from their surfaces, and inclined to the longitudinal course in the intermediate space. This structure, Mr. Owen states, proves that the vertebra cannot belong to the *Poikilopleuron Bucklandi*.

A caudal vertebra also from Buckingham, and assigned by Professor Owen to the middle part of the tail, on account of the development of short, narrow transverse processes just below the neurapophysial sutures, exhibited a centrum of a subtriangular form, with one angle inferiorly and the other two at the origin of the transverse processes, but all three largely rounded off. The marginal circumference of the centrum was convex, and separated from the lateral or free surface by a rough, irregular, elevated ridge, the inferior part of which encroached upon the under surface of the vertebra in the form of two semicircular facets, both anteriorly and posteriorly. The free surface of the vertebral centre is marked by the coarse lines of the bony fibrous structure, decussating like an irregular net-work. The spinal canal of this specimen did not sink into the body of the vertebra. The size of this vertebra, and the proportions and position of neurapophyses and hæmapophysial articulations, might suggest a relationship of the animal to which it belonged with the Cetacea ; but it differs, Mr. Owen says, in the concavity of the terminal articulations, which show no sign of separation as laminar epiphyses, and more particularly in the place of

the origin of the transverse process being close to the neurapophysis instead of proceeding from the middle of the side of the centrum. In these deviations from the Cetacea, the *Cetiosaurus* approaches, the author states, the saurian order.

Mr. Owen then describes, with his wonted minuteness and perfect acquaintance with the subject, other caudal vertebræ found at Blisworth, but it is not possible to abridge the details.

Among the remains discovered near Chipping Norton are eleven caudal vertebræ without transverse processes, and therefore assigned by the author to the terminal half of the tail. They progressively diminished in transverse diameter from five inches to two inches, but without losing in equal ratio their length, which continues the same, or five and a half inches in the vertebra which has only three inches and three lines of breadth, five inches in that which is two inches and nine lines broad, and four inches in that which has a breadth of two inches. These eleven vertebræ do not constitute, Mr. Owen shows, a regular sequence, but detached links of the termination of the spinal column. In all the existing genera of Cetacea the posterior caudal vertebræ become shorter in proportion to their thickness, and the terminal ones are depressed. The slender elongated form of the corresponding vertebræ in the *Cetiosaurus*, is, Mr. Owen shows, a striking crocodilian character; and he adds, it is important to observe that not any of the series of caudal vertebræ described in this paper exhibit the vertical canals or perforations of the side of the centrum or base of the transverse process which so peculiarly characterizes most of the cetacean caudal vertebræ.

In his comparison between the vertebræ of the *Cetiosaurus* and the *Poikilopleuron*, Professor Owen states that the caudal vertebræ of the former resemble those of the latter and most other reptiles from strata below the chalk in the articular surfaces being slightly concave; and the vertebræ of the *Poikilopleuron*, especially in the elongated and rounded form of the body; in its median compression, and in the articulation of the hæmapophyses to the inferior part of the vertebral interspaces, though they are larger; on the contrary, the *Cetiosaurus* vertebræ differ in their proportions, in their structure, as in the absence of the remarkable medullary cavity in the middle part of the centrum of the *Poikilopleuron*; in the shortness of the neurapophyses as compared with the centrum; and in other minor points, which are fully detailed by Professor Owen.

The author then proceeds to institute further comparisons between the vertebræ of the *Cetiosaurus* and other reptilia: thus he shows that they differ from the vertebræ of the Crocodilians in retaining the cylindrical form of the body to the end of the tail, instead of being compressed and four-sided; that there is no trace of the vertical median division which the bodies of the caudal vertebræ present in *Iguanæ*, *Anolides* and other Lacertians; that they are not only larger than in the *Megalosaurus*, but relatively longer; that they differ from the anterior caudal vertebræ of the *Iguanodon*, which are nearly as large, in the absence of the well-marked concavity below the transverse processes, in the form of the centrum not being so quadrilateral, and especially in the transverse breadth

of the inferior surface being less; and from the posterior caudal vertebræ of the Iguanodon, which slightly increases in length, in being less compressed and the centrum not having a triangular form; the slender terminal caudal vertebræ of the Iguanodon are also hexagonal, and not cylindrical as in the Cetiosaurus.

As there is no known extinct saurian which can so nearly compete in size with the Cetiosaurus as the Iguanodon, it is fortunate, Prof. Owen observes, that the distinguishing characters are so well marked and easily recognizable.

*Dorsal vertebra.*—The only portion of a dorsal vertebra described in the memoir is the extremity of a spinous process, the posterior surface of which is rough and flattened, 4 inches across, at about the same distance below the end of the spine; the sides are traversed to a certain extent by a longitudinal ridge, anterior to which they are concave and smooth, but their anterior margin is again flattened and rough, though it is not so broad as the posterior.

In referring all the vertebræ described in this paper to the same species of saurian, Prof. Owen admits that they present a somewhat greater variety of form and proportion in different regions of the tail than is observable in that part of the vertebral column in the smaller and recent species of Crocodile or Lizard; not only becoming larger in proportion to their thickness, but increasing slightly in length for a short distance as they recede from the sacrum. They appear likewise to exchange from a cylindrical to a subtriangular form of the body, but to resume the cylindrical shape in the terminal half of the tail. These modifications, he says, are possible, as in the *Plesiosaurus brachydeirus* still greater discrepancies in the proportions of the vertebræ prevail; and they are inferior in degree to any of the modifications which distinguish the vertebræ of known genera of saurians from those under consideration, in pointing at their distinguishing features from the hitherto known sauria; and in thus treating of them collectively, the inference that they belong to the same gigantic species is, the author observes, almost irresistible, that they belong to a new and distinct genus, which, on account of the vertebræ approximating in size and structure to the vertebræ of the whale, he has termed Cetiosaurus.

In the cuttings for the London and Birmingham Railway near Blisworth, there were found, scattered over an area of 12 feet by 8 feet, the following remains:—1. A bone resembling the episternal of an Ichthyosaurus, the length or antero-posterior extent of the preserved portion of the median plate being  $1\frac{1}{2}$  foot, and the breadth of the posterior fractured end 5 inches, from which it gradually expands to the root of the side branches, where its breadth is 1 foot. From its obtuse termination to the end of the longest branch is  $2\frac{1}{2}$  feet, and from this end to that of the opposite branch  $4\frac{1}{2}$  feet. 2. The remains of a coracoid and scapula apparatus of equally gigantic proportions. 3. A fragment, considered to be the shaft of a humerus, 1 foot 9 inches in length, 6 inches in diameter across the middle and 8 inches across the widest end. 4. A portion of the opposite humerus. 5. Another fragment, believed to be part of a radius or ulna, about a yard in length, 6 inches across the proximal end,

and 5 inches across the middle of the shaft. 6. A slightly curved portion of a rib, a yard long and from  $1\frac{1}{2}$  to 2 inches thick. 7. Five caudal vertebrae agreeing in dimensions with the vertebrae of Chipping Norton.

Numerous fragments of long bones without a trace of a medullary cavity have been found at Chipping Norton, and correspond in magnitude with the vertebrae. The articular surfaces which are preserved are covered with large tubercles for the attachment of thick cartilages. The best preserved fragments are considered to belong to metacarpal or metatarsal and phalangeal bones, and are therefore, Prof. Owen says, decisive evidence against the cetacean nature of the animal; but he adds, they possess characters by which they may be distinguished from the corresponding bones of known extinct gigantic saurians. One of these bones, believed to be a metacarpal or a metatarsal, is double the bulk of the largest analogous bone of a full-grown elephant, though the metacarpals or metatarsals are much smaller in proportion in Saurians than in Pachyderms. The bone is 7 inches in length, 9 in circumference in its middle, 5 in the antero-posterior diameter of its proximal end, and 4 inches 8 lines in the transverse diameter of the distal end. A proximal phalanx is shown to be remarkable for its short and broad proportions, which are more massive than those of the phalanges of existing Crocodilians or of the Poikilopleuron.

An ungual phalanx, also found at Chipping Norton, was 6 inches in length,  $2\frac{1}{2}$  in breadth, and upwards of 3 in depth. It was slightly curved, obliquely compressed, obtusely terminated with a shallow, concave, trochlear articular surface, divided by a vertical convexity; it was marked on each side by a smooth curved groove, 3 inches in length, with the concavity downwards, and the lower edge projecting beyond the upper at the posterior part of the groove; but it is shown to be by no means produced in so large and thick a ridge as that which characterizes each side of the more depressed and broader phalanx of the Iguanodon. From the ungual phalanges of that Saurian it differs in being much less compressed from side to side and less curved downwards. It vastly surpasses in size any of the ungual phalanges of the Poikilopleuron. A smaller ungual phalanx, resembling in general shape the above, was found at Chipping Norton; and portions of metacarpal or metatarsal bones, agreeing in form and size with the fragments obtained at Chipping Norton, have been discovered at Buckingham: also a fragment 8 inches long, which Prof. Owen considers to have belonged to a radius, a fibula, or a long distal phalanx.

With reference to a comparison of the remains of the Cetiosaurus with those of the Polyptychodon, the bones of the extremities present in both cases the cancellous structure throughout the central part, which indicates aquatic rather than terrestrial habits. Prof. Owen states that he has not found any of the remains of the extremities of the Cetiosaurus to agree exactly in shape with those belonging to the Polyptychodon; also that no specimen of a tooth agreeing in characters with the teeth of the Polyptychodon has been detected in secondary strata inferior to the greensand. Certain

large conical teeth, found in the Malton oolite, may, Mr. Owen thinks, appertain to the *Cetiosaurus*, but he is of opinion that they more probably belong to the *Steneosaurus*.

In conclusion, it is stated that the vertebræ described in the paper prove the existence of a saurian genus distinct from the *Megalosaurus*, *Steneosaurus*, *Poikilopleuron*, *Plesiosaurus*, or any other large extinct reptile, remains of which have been discovered in the oolitic series; that the vertebræ, as well as the bones of the extremities, prove its marine habits; and that the surpassing bulk and strength of the *Cetiosaurus* were probably assigned to it with carnivorous habits, that it might keep in check the *Crocodylians* and *Plesiosaurs*.

5. "On the age of the Tertiary beds of the Tagus, with a Catalogue of the Fossils," by James Smith, Esq., of Jordan Hill, F.G.S.

During a visit to Portugal in 1840, Mr. Smith made a collection of the organic remains in the tertiary deposits near Lisbon, for the purpose of ascertaining their relative geological age. Since his return to England he has carefully examined the collection, assisted by Mr. George Sowerby, and ascertained that the series of beds from which they were obtained belong to the miocene division of the tertiary system, and to that portion of it which includes the Bordeaux and Dax beds, rather than to any other yet described deposit. He has, however, determined, by a careful comparison of the Lisbon fossils with those given in the works of MM. de Basterot and Grateloup, and with his own collection of Bordeaux organic remains, that there is a greater difference than can be ascribed to geographical distance alone; but he hesitates to assign to the Lisbon beds either a more ancient or a less ancient date. The proportion of recent shells, he states, affords no assistance, as, according to M. de Basterot, the existing species in the Bordeaux basin equal 23 per cent., and according to M. Grateloup, 37 per cent., whilst Mr. Smith's collection of Lisbon fossils gives 28 per cent. The author is fully convinced of the soundness of the principle of determining the comparative age of a tertiary deposit by the proportion of recent species; but he is of opinion, on account of the great difficulty of defining species, that it is only possible to arrive at an approximation sufficiently near to decide to which of the great divisions of the tertiary system a set of beds may belong, and not to the precise relative antiquity of two deposits of nearly the same age.

Prof. Agassiz has decided that several of the new species of Lisbon shells occur in the molasse of Switzerland, and he considers the two series of strata as nearly contemporaneous.

Mr. Smith refers to Mr. D. Sharpe's memoir on the neighbourhood of Lisbon\*, for a description of the mineral structure of the formation, confining his own remarks to pointing out the localities and position in the series from which the fossils were obtained.

\* Proceedings, Geol. Soc., vol. iii. p. 28, 1839; also Geol. Trans., Second Series, vol. vi. p. 1. A list of tertiary shells is given in p. 113.



Date: Fri, 9 Mar 2007 02:44:36 -0800 (PST)  
From: pathman@sunsite4.berkeley.edu  
To: nrlfreq@library.berkeley.edu  
Subject: UCB Library Pathfinder NRLF Request/WEB DELIVERY  
X-Mailer: WebZMailer

\*\*\*\* For email communications contact nrlfreq@library.berkeley.edu \*\*\*\*

An NRLF request was made by the following patron through Pathfinder

WEDEL, MATHEW JOHN  
ID: 15633183  
Status: 2  
Dept: INTEGRATIVE BIOLOG  
Email: sauropod@berkeley.edu

Volume: 3 Date: 1841

Article Title: A description of a portion of the skeleton of Cetiosaurus

Article Author: Owen, R

Pages: 457-462

Barcode:

Loc/Call No: Earth Sci QE1 .G49 Shelved at NRLF: 1826/27-1844/45, 1952/53-1966/67,  
1968/69-1970/71

Title: Proceedings.

Gladis Number: GLAD16933913

NRLF will notify you at sauropod@berkeley.edu when the article you requested is available on the web (normally within two working days), or if there is a problem.

BA 168 257

~~DTD~~ Ariel  
NRLF Access Services  
Date 3/9/07  
Initials ALP  
PS 7000A  
Pages 7

UCB.09K